

# Spatial variation in vegetation height as an indicator of aboveground carbon stocks in grazed grasslands


Klumpp K., Darsonville O., Bloor J.M.G.

INRAE, University of Clermont Auvergne, VetAgro Sup, UMR UREP, Clermont-Ferrand, France



## Background

Spatial heterogeneity in plant and soil properties is known to influence ecosystem functions, but the linkages between spatial variation and ecosystem services in grasslands are unclear. **Here we examine within-field variation in sward structure (vegetation height) and test whether indices of spatial heterogeneity can be used as a simple indicator of aboveground production services or carbon stocks over time in upland grasslands.**



Left : Study location in the Massif Central region of France ; Right : On-site grazing.




Grid measurements at the study site. Photos: C.Schmidt

## Experimental approach

- Measurements were carried over 6 years at a **long-term research platform** in France (SOERE-ACBB: Laqueuille) on **two grazed grasslands** (high stocking rate and N inputs, intensive; low stocking rate and no N inputs, extensive).
- **Field-scale standing biomass** and potential productivity was assessed across each paddock at five dates per year during the growing season.
- Maximum height of both green and senescent shoots were determined at the end of each grazing season on **30m x 20m grids** using a sward stick (117 points per grid, 2.5m distances). Standard deviation (SD) and coefficients of variation (CV) were used to assess the variation in plant height.

## Results

- Mean annual productivity was greater in the 'Intensive' treatment compared to the 'Extensive' grazing treatment ( $F_{1,11} = 54.5$ ,  $P < 0.001$ ), but standing biomass showed no significant difference between grazing treatments during the study (Figure 1A/B).
- Absolute variation in **green vegetation height** at the end of the growing season (green height SD) showed a **positive relationship** with mean annual **field-scale standing biomass** in both grazing treatments (Figure 1C).
- Metrics of **green height** were **unrelated to field-level productivity**, and no significant relationship was found between metrics of variation in senescent height and either field-level productivity or standing biomass ( $P > 0.05$ ).



**Fig. 1.** Interannual variation in field-scale annual biomass production (A) and mean standing biomass (B); data are means  $\pm$  SE. Relationship between within-field variation in green vegetation height (standard deviation) and estimations of mean annual field-scale standing biomass during the study (C). Grazing treatments are given by: Intensive, orange-filled bars/circles; Extensive, blue-filled bars/triangles.

## Take-home Message

Our results suggest that within-field variation in end-of-season **green vegetation height** may provide an integrated **indicator of biomass state** during the year (i.e. quantity of standing biomass available for ingestion over time), with implications for the estimation of **aboveground carbon stocks and carbon input into the soil**. However, simple metrics of within-field variation do not appear to provide useful proxies of biomass fluxes in this grassland system.